Original Research Article



Saudi Medical Journal for Students (SMJS)

Official Journal of Faculty of Medicine University of Tabuk ISSN: 1658-8274 (Print version); 1658-8282 (Electronic version)

THE EFFECT OF PARENTAL STRESS ON CHILDREN'S PSYCHOLOGY DURING COVID-19 CRISIS IN MAKKAH REGION: CROSS-SECTIONAL STUDY

Murouj Almaghrabi¹, Ahdab Bawashkhah¹, Arwa Ahmed Abba¹, Ghady Algethami¹, Maram Alshareef²

¹Medical student, College of Medicine at Umm Al-Qura University, Al-Abdia Main Campus, Makkah, Saudi Arabia ² Assistant Professor, Department of Community Medicine and Pilgrims Health Care, Faculty of Medicine, Umm Al-Qura University, Makkah, Saudi Arabia

***Corresponding author:** Murouj Almaghrabi, College of Medicine, Umm Al-Qura University, Makkah, Saudi Arabia. Tel: (+966) 536251063.ORCID number: https://orcid.org/0000-0001-7357-9269. Email: murouj@outlook.com, <u>437000844@st.uqu.edu.sa</u>

ABSTRACT

Background and aim: The rapid transmission of COVID-19 has resulted in multiple restricted precautions that may lead to psychological impacts on individuals. Hence, it is essential to identify mental health issues to find the best solutions. Therefore, our study aims to inspect the effects of COVID-19 on children's mental health concerning the parents' stress levels in the Makkah region. **Methods:** A descriptive anonymous cross-sectional study was conducted via a modified webbased questionnaire to assess children's psychological impact in correlation with parental stress during the coronavirus pandemic in the Makkah region, Saudi Arabia.

Results: This study includes 845 participants represented by 339 parents. Changes in the emotional state and behaviors among children were found in a total of 138 children (27.06%), in which the highest age group was 5-7 years (42.8%; 59). Moderate and high levels of stress were detected in parents (68.1% and 4.7%, respectively). No significant relation was seen between parental stress and psychological changes among their children. However, statistical significance was found between changes in children's psychology and family history of COVID-19 infection (P=0.001). Parents' nationality showed a statical significance difference for moderate to a high level of stress among foreigner parents (P=0.039 and 0.043, respectively, for mothers and fathers). **Conclusion:** The current study revealed no significant relationship between parental stress and psychological impact on their children. However, higher levels of parental stress, as well as a significant change in children's emotional state and behaviors, were seen as an impact of COVID-19 quarantine.

Keywords: Children, psychology, Parental-stress, Quarantine, COVID-19, Makkah.

To cite this article: Almaghrabi M, Bawashkhah A, Abba AA, Algethami G, Alshareef M. Saudi Medical Journal For Students: The Effect of Parental Stress on Children's Psychology during Covid-19 Crisis in Makkah Region: Cross-Sectional Study 2022;3(1):63-76

INTRODUCTION

Coronavirus disease 2019 (COVID-19) is an emerging infectious disease from the family of coronaviruses. It was initially discovered in China, Wuhan. [1,2] By November 8, 2020, World Health Organization (WHO) had registered more than 49 million confirmed cases of COVID-19 and more than 1 million deaths worldwide. [3] Due to its rapid transmission among people, many precautions have been taken worldwide to limit its spread, one of which is home quarantine. [4,5]

Besides the benefits of quarantine, it is stated that the endless time, the fear of getting infected and spreading it to others, the routine, and the lack of supplies such as food, clothes, and regular medical care are all considered stressors of quarantine that lead to frustration, anxiety, and anger. [6] Some ways to overcome the impact of quarantine include; giving adequate information about the length of quarantine, minimizing its duration, and raising awareness about the importance of healthy communication as it reduces the feeling of isolation and stress. [6]

Throughout the COVID-19 crisis, parents observed changes in their children's state of mind and attitude during the quarantine. The most common was the inability to concentrate, boredom, nervousness, desolation, and worries. [7] Parents can observe stress among children through signs like crying, sadness, extreme worry, unhealthy habit, headache or pain from an

unknown cause, and more. [8] Accordingly, parents can work effectively with their children during COVID-19 to minimize their psychological impact [9]. This could be done through different methods such as increasing communication with them to know their fears and answer their questions [8,10]. encouraging them to do some collaborative activities, make an active routine for them, relax, eat well, exercise, and get plenty of sleep. [8,10,11] In addition, communication with family and friends through online applications can also help the child. Parents need to know the signs of stress to deal with it as early as possible. [8,10]

Unfortunately, parents' stress can be a barrier, especially in a period of crisis like COVID-19. It could affect children's stress and stress expressions. [12] It has been proved that COVID-19 and quarantine have affected the population's health by increasing stress and depression. [13] A previous study examined the emotional impact of quarantine on children and adolescents. They found that behavioral and mental state changes were noticed in order from highest to lowest difficulty concentrating, boredom, and irritability. As expected, the children were exposed to screens more, as they spent less time doing physical activity and more time sleeping. [7] Another study concluded that the stress resulting from the COVID-19 outbreak harmed the parents' behavior towards their children. [14] To the best of the authors' knowledge, no study has covered the impact of quarantine on children's mental

state and its relation to the level of their parents' stress. Hence, it is important to investigate this relation as it affects the future community health and well-being, as well as to help to identify the effect of parent stress on children's psychology for further action regarding psychological rehabilitation programs, so we are conducting this study. The study aims to inspect the impact of the COVID-19 outbreak on children's mental health concerning the parents' stress levels in the Makkah region.

METHODS

Ethical approval and consent

Ethical approval was obtained from the Internal Review Board (IRB) at Umm Al-Qura University (UQU), Makkah, Saudi Arabia. Approval No. (HAPO-02-K-012-2020-06-409). An electronic informed consent was obtained from each participant prior to answering the questionnaire. Confidentiality was assured. Names or phone numbers were not requested from any participant.

Research design, population, and sample size determination

This descriptive cross-sectional study was conducted among parents and children in the Makkah region, Saudi Arabia. A sample of 845 individuals with 339 parents and 506 children was collected randomly through a self-administered online questionnaire from June 25 to July 8, 2020. Due to limited census information and publications of children and parents in the Makkah region, which decreased the accuracy of sample size determination, we considered all residents in Makkah to be our population size. The

required sample size was calculated by OpenEpi version 3.01, [15] in consideration of the following: the population size is approximately 2,042,000 inhabitants (as reported in the last statistics for this year 2020), keeping the confidence interval (CI) level at 95% and considering Anticipated % of frequency as 50% and taking design effect as 1. The sample size was calculated to be 385 participants. Our study targeted residents of the Makkah region, parents of all ages with at least one child, and children. We excluded any child over 15 years as they considered beyond the standard age of children. In addition, we excluded any child less than three years as the tool will not help explain their impressions. Finally, special needs children were also excluded from the study. A computer-assisted Google survey program was used for the questionnaire, as face-toface contact was not allowed during that period.

Questionnaire structure

A modified web-based questionnaire was used to assess the children's psychological impact along with parental stress during the coronavirus pandemic. It is distributed bilingually (Arabic and English) via different platforms social on media. Arabic questionnaire was translated back to English for analysis and publication. This questionnaire was inspired, combined, and modified using two different sources, and all modifications were done after permission. [7,16] Our adapted questionnaire consists of three sections, as follows:

Section 1. Socio-demographics and family characteristics: included six questions about whether they are living with their child or not,

age, nationality, educational level, marital status, and occupation. As well as seven questions about residential area, number of family members in the house, family income, description of the house (whether have outdoor space or not), house space (number of rooms), place where the child lived during quarantine, and any family member with a previous history of covid-19 infection.

Section 2. Parental perception of the emotional and behavioral effects of the quarantine on their child/children; this included 30 factors of different behavioral and emotional changes; each factor consisted of 5 options ranging from "much less compared to before quarantine" to "much more compared to before quarantine."

Section 3. Perceived mental stress level among parents is a classic stress assessment instrument that helps to understand how different situations affect feelings and perceived stress. It included ten questions about parents' feelings and thoughts during the last month. For each question, we used the 5-point Likert scale for the participant to admit their agreement level. These five points were: ("never," "almost never," "sometimes," "fairly often," and "very often").

Statistical analysis

Participants' data were entered automatically via an Excel sheet where it was cleaned, coded, and fed to statically software IBM Statistical Package for the Social Sciences (SPSS) version 23 for conducting all statistical issues. Descriptive statistics were used (frequency and percentage); statistical significance was determined at $p \le 0.05$. And CI was considered as 95%. The Chi-square test was applied for categorical data analysis. Additionally, Pearson's Correlation Coefficient (PCC) was run to discover the relation between parental stress and their children's psychological impact. A total of 339 parents registered their family demographic information, including their own and partner's demographics and some general information about the family.

Then, each parent registered their perception of quarantine's emotional and behavioral effects on their child/children, who reached 506 children. Finally, they individually logged their responses to the Perceived Stress Score (PSS).

RESULTS

Socio-demographic characteristics of the participants

The demographical data of 845 participants (parents/ children) of the study were represented by 339 parents, as demonstrated in Table1 . Most respondents were mothers (75.8%) and lived in Makkah (76.7%). Most participants had 5-7 family members in the house (52.5%), followed by 2-4 members (36.3%). It was noticed that most participants did not own outdoor space at home (63.1%), while those who owned outdoor space, either garden or balcony, were 125 (36.9%). Family income was over 10,000 SAR in 176 participants (51.9%), while only 6 (1.8%) had no income. Regarding the living status information, most children stayed with their parents (65.8%). The results showed only 85 (25.1%) had got infected with COVID-19 in the family. The relative relation of 37 of them was the first degree. Regarding parental data, almost all of them were living with their children (98.5% and 90.3% for mothers and

SMJS is the official journal of the Faculty of Medicine, University of Tabuk. All rights reserved with SMJS. © SMJS 2021

General	l informati	on of the family (n=339)				
C	haracteristi	с		No (%)		
Parent		Mothers		272 (75.8%)		
		Fathers		82 (24.2%)		
Residence area		Makkah	260 (76.7%)			
		Jeddah	79 (23.3%)			
Number of family member in the house		2-4	123 (36.3%)			
		5-7		178 (52.5%)		
		8-10		34 (10%)		
		More than 10		4 (1.2%)		
Description of the house		Outdoor space (garden)		72 (21.2%)		
		Outdoor space (balcony)		53 (15.6%)		
		No outdoor space		214 (63.1%)		
House space (number of rooms)		1-2 room/s		6 (1.8%)		
		3-4 rooms		130 (38.3%)		
		5 rooms		103 (30.4%)		
		More than 5 rooms	100 (29.5%)			
Family income		< 5000 SAR		41 (12.1%)		
5		5000 - 10,000 SAR		116 (34.2%)		
		> 10,000 SAR		176 (51.9%)		
		No Income		6 (1.8 %)		
During the quarantine, where did the child/childre	en lived?	With their parents only		223 (65.8%)		
		In the big family house		99 (29.2%)		
		With some family member	r (uncle or aunt)	17 (5%)		
Was anyone in the family diagnosed with covid-1	9?	Yes		85 (25.1%)		
, , , ,		No	254 (74.9%)			
If yes, what is the relative relation?		The child		3 (0.9%)		
This is a specific question for those who answer	ed 'ves' in	First degree		37 (10.9%)		
the previous item $(n=85)$	5	Second degree	28 (8.3%)			
		Distant family member or	17 (5%)			
				254 (74.9%)		
Pa	rents' chai	acteristics (n=339)				
Characteristi	ic		Mothers	Fathers		
Does the parent live with the child /children?	Yes		334 (98.5%)	306 (90.3%)		
1	No		3 (0.9%)	23 (6.8%)		
	Passed a	wav	2 (0.6%)	10 (2.9%)		
Age	<29		63 (18.7%)	19 (5.8%)		
	30-39		121(35.9%)	110 (33.4%)		

Table1 : Socio-demographics and sample characteristics of the participant

Pa	rents' characteristics (n=339))	
Characterist	ic	Mothers	Fathers
Does the parent live with the child /children?	Yes	334 (98.5%)	306 (90.3%)
	No	3 (0.9%)	23 (6.8%)
	Passed away	2 (0.6%)	10 (2.9%)
Age	≤29	63 (18.7%)	19 (5.8%)
	30-39	121(35.9%)	110 (33.4%)
	40-49	126 (37.4%)	112 (34%)
	≥50	27 (8%)	88 (26.7%)
Nationality	Saudi	293 (86.9%)	290 (88.1%)
	Non-Saudi	44 (13.1%)	39 (11.9%)
Educational level	Less than high school	24 (7.1%)	40 (12.2%)
	High school/ diploma	72 (21.4%)	93 (28.3%)
	Collage or above	241 (71.5%)	196 (59.65%)
Marital status	Married	312 (92.6%)	323 (98.2%)
	Divorced/ Widow	25 (7.4%)	6 (1.8%)
Occupation	Health care provider	23 (6.8%)	26 (7.9%)
	Retired	13 (3.9%)	35 (10.6%)
	Field work	69 (20.5%)	107 (32.5%)
	Office work	72 (21.4%)	137 (41.6%)
	Not working/ Student	160 (47.4%)	24 (7.3%)
fathers respectively). Parents' ages s remarkable convergence, as most m	show a ages are nothers' followed l	between 40-49 ov 30-39 years (3	years (37.4%), 35.9%). On the

SMJS is the official journal of the Faculty of Medicine, University of Tabuk. All rights reserved with SMJS. © SMJS 2021

other side, most of the fathers' age were between 40-49 years in 112 (34%), followed by 30-39 years in 110 (33.4%) participants. Most of the parents were Saudis (86.9%, 88.1%), married (92.6%, 98.2%), and their educational level at college or above (71.5%, 59.65%) for both mothers and fathers, respectively. Most mothers were not working/students (47.4%), while most fathers had office work jobs (41.6%).

Parental perception of the emotional and behavioral effects of the quarantine on their child/children

One of the main issues in our study is to inspect the psychological effect of quarantine on children. Thus, 30 factors (previously prescribed in methods-section 2.2) were listed in a multiple-choice grid table to accomplish the objective. One point was given to each participant who chose one of the following options "sometimes, fairly often, or very often." For each participant, a maximum score of 30 was calculated. Furthermore, a score of 16 or above was a significant psychological impact. Table 2 represents the total number of results, in which 138 children (27.06%) have recorded a significant change in their psychology (emotional state and behavior changes). This number is distributed from the highest to lowest between three groups: children aged 5-7 years (42.8%), 8-10 years (36.2%), and 11-15 years (21.0%). On the other hand, 368 children (72.93%) of the total number of children (506) have a negative psychological impact.

Prevalence of perceived mental stress level among parents and gender differential

Error! Reference source not found. p rovides a total score of stress among parents. Nearly half of the patricians have moderate stress (68.1%), including fathers and mothers. Followed by low-stress levels detected in 92 (27.1%) participants. Lastly, high levels of stress were seen in 16 (4.7%) parents. No significance was detected between different groups of genders.

Association between parental stress and the psychological impact on their children during the quarantine

Pearson's Correlation Coefficient (PCC) was run to discover the relation between variables. It revealed no significant relationship between parental stress and psychological impact on children.

Family characteristics in association with parental stress and the psychological impact of quarantine on children

Error! Reference source not found. r epresents the distribution of family characteristics in association with parental stress and psychological impact on their children. We found that in children with psychological changes, 37.7% of them had a family member with a previous family history of COVID-19 infection compared to 62.3% of children with no previous family with recorded statistical history, а significance (P=0.001). Other factors including residence area, number of a family member in the house, family income, description, and space of the house, as well as the place where the child lived during quarantine, had no significant relation with psychological changes among children (p=

SMJS is the official journal of the Faculty of Medicine, University of Tabuk. All rights reserved with SMJS. © SMJS 2021

	Total (n=506)		Childre	Children between		Children between		Children between	
			5-7 years		8-10 years		11-15 years		
	(n=193)		(n=156)		(n=157)				
	Ν	%	Ν	%	Ν	%	Ν	%	
Significant changes in	138	27.06%	59	42.8%	50	36.2%	29	21.0%	
children's psychology									
No significant psychological	368	72.93%	134	36.4%	106	28.8%	128	34.8%	
impact									

Table 2: Parental perception of the emotional and behavioral effects of the quarantine on their child/children (n=506)

 Table 3 Perceived mental stress level among parents and gender differential (n=339)

PSS score category	Total (n=339)		Mothe	Mothers (n=272)		Fathers (n=82)		
	Ν	%	Ν	%	Ν	%		
Low stress	92	27.1%	68	26.5%	24	29.3%		
Moderate stress	231	68.1%	176	68.5%	55	67.1%	0.795	
High stress	16	4.7%	13	5.1%	3	3.7%		

Notes: *Pearson chi-square test. The difference is significant at P value ≤0.05

0.114, 0.172, 0.301, 0.213, 0.078 and 0.634, respectively).

Regarding parents with moderate to high stress levels, 63.8% lived in Makkah during quarantine time, while 36.2% the participants lived in Jeddah, with statistical significance (P=0.002). As for the number of family members in the house, 55.2% of fathers and 51% of mothers with moderate to high stress levels have 5-7 members compared to 3.4%, and 1.1% of parents have than ten members (P=0.446). more Regarding family income among parents with moderate to high levels of stress, 60.3% of fathers and 48.1% of mothers have > 10,000 SAR income, while 0.0% of fathers and 2.1% of mothers had no income (P=0.097). Additionally, no statistical significance was recorded in the house space, where the child lived during quarantine, nor previous family history of COVID-19 infection (P=0.413, 0.478, and 0.136, respectively).

Parental demographics in association with their level of stress during the quarantine

The distribution of parental stress during the quarantine of the COVID-19 pandemic by their socio-demographic data is shown in Table 5. Exactly 98.4% of mothers living with their child/children had moderate to a high stress level compared to mothers who were not living with their child/children (1.6%) with recorded statistical significance (P=0.024). Also, 81% of mothers aged 40-49 years had moderate to high-stress levels compared to 7.9% of mothers aged more than or equal to 50 years (P=0.016). As for nationality, 90.5% of mothers were Saudis and had moderate to high-stress levels compared to non-Saudis mothers (9.5%)

SMJS is the official journal of the Faculty of Medicine, University of Tabuk. All rights reserved with SMJS. © SMJS 2021

(P=0.039). Other including factors. educational level, marital status, and occupation, had no significant relation to stress. On the other side, Saudi fathers had moderate to high-stress levels (81%) compared to 19% of non-Saudi fathers, with recorded statistical significance (P=0.043). Other factors, including living with their child, age, educational level, marital status, and occupation, had no significant relation to stress level (P=0.180, 0.111, 0.120, 0.445, and 0.586, respectively).

DISCUSSION

This study is among the first studies done in Saudi Arabia to examine the psychological impact of COVID-19 quarantine on children and their parents and discover the relationship between them. The setting of our study was chosen to be Makkah due to its higher numbers of COVID-19 recorded cases compared to other cities in the kingdom. [17] As well as special governmental precatory measures that have been done for this region only, such as temporary suspension of Muslim entry to perform the Umrah. All these changes and new situations may affect the emotional and behavioral well-being of the individuals.

Children's psychological impact

Our study revealed that from a total of 506 children, significant changes in psychology were detected in 138 (27.06%) participants. The highest age group was the younger children between 5-7 years (42.8%). These results are in agreement with a preliminary study conducted in China during the COVID-19 epidemic, which reported that children aged 3-6 years were more inclined to display

symptoms than their peers from older age groups (P =.002). Examples of these symptoms include clinginess and afraid of family members getting the infection. [10] Anyhow, multiple studies have also reported changes in children's emotional state and behaviors during the quarantine of COVID-19. A previous study in Italy and Spain revealed that 85.7% of the parents had noticed these changes among their children. most frequent symptoms The were. respectively. difficulty in concentrating (76.6%), boredom (52%), and irritability (39%). [7] Several studies have also praised the negative effect of quarantine on children's psychology [6,18-20]. According to these terrifying results, there is an urgent need to detect these changes among children as early as possible to solve them much easier. In such situations, parents play an essential role, as the key to identifying psychological issues and comforting children in prolonged isolation is close and open communication with the children. [21]

Parental perceived stress

Assessment of parents' psychological health is essential for parents and their children. Consequently, an internationally validated tool for the assessment of perceived stress was used in this study. [16] Our results showed that more than half of the parents had a moderate stress level (68.1%). Moreover, 4.7% of parents had a high level of stress. These results are consistent with several studies that have examined stress or other psychological impacts among parents during various pandemics, including the COVID-19 crisis. These studies were done all over the globe, including in the United States [22,23],

SMJS is the official journal of the Faculty of Medicine, University of Tabuk. All rights reserved with SMJS. © SMJS 2021

Variable	Total number of		Moderate/high	Moderate/high	Р	
	children with +ve	value	stressed mothers	stressed fathers	value	
	psychological impact		(n=189)	(n=58)		
	(n=138)					
Residence area						
Makkah	112 (81.2%)	0.114	158 (83.6%)	37 (63.8%)	0.002*	
Jeddah	26 (18.8%)	0.114	31 (16.4%)	21 (36.2%)	0.002*	
Number of family member in the	he house					
2-4	31 (22.5%)		68 (36%)	20 (34.5%)		
5-7	88 (63.8%)	0 172	97 (51.3%)	32 (55.2%)	0.446	
8-10	18 (13%)	0.172	22 (11.6%)	4 (6.9%)		
More than 10	1 (0.7%)		2 (1.1%)	2 (3.4%)		
Family income						
< 5000 SAR	17 (12.3%)		30 (15.9%)	3 (5.2%)		
5000 - 10,000 SAR	41 (29.7%)	0.201	64 (33.9%)	20 (34.5%)	0.007	
> 10,000 SAR	74 (53.6%)	0.501	91 (48.1%)	35 (60.3%)	0.097	
No Income	6 (4.3%)		4 (2.1%)	0		
Description of the house						
Outdoor space (garden)	24 (17.4%)		38 (20.1%)	17 (29.3%)		
Outdoor space (balcony)	24 (17.4%)	0.213	31 (16.4%)	9 (15.5%)	0.331	
No outdoor space	90 (65.2%)		120 (63.5%)	32 (55.2%)		
House space (number of rooms)					
1-2 room/s	2 (1.4%)		4 (2.1%)	0		
3-4 rooms	62 (44.9%)	0.079	67 (35.4%)	16 (27.6%)	0.412	
5 rooms	43 (31.2%)	0.078	60 (31.7%)	20 (34.5%)	0.415	
More than 5 rooms	31(22.5%)		58 (30.7%)	22 (37.9%)		
The place where the child/child	lren lived in during quara	antine				
With their parents only	97 (70.3%)		124 (65.6%)	43 (74.1%)		
In the big family house	35 (25.4%)	0.634	56 (29.6%)	13 (22.4%)	0.478	
With some family member	6 (4.3%)		9 (4.8%)	2 (3.4%)		
Any family member with a pre-	vious history of covid-19	infection				
Yes	52 (37.7%)	0.001*	54 (28.6%)	7 (12.1%)	0.136	
No	86 (62.3%)	0.001	135 (71.4%)	51 (87.9%)	0.150	

Table 4: Family characteristics in association with parental stress and psychological impact among children during quarantine of COVID-19 pandemic (n=385)

* Statical significance p-value was detected at $(p \le 0.05)$ using Pearson chi-square test.

China [24,25], Italy [26], and Korea. [27] These results make sense, as individuals face many difficult challenges during such a crisis as the COVID-19 pandemic. Copping with these struggles is a keystone. A study published in May 2020 reported that psychosocial crisis management and intervention models should be urgently implemented by the government or health care professionals to better cope with these psychosocial problems from various strata of society. The use of internet infrastructure, technology, and social media to combat both pandemic and info-demic needs to be introduced. [28]

Relation between parental stress and psychological impact on their children The results of the present study support that there was no relation between parental stress

SMJS is the official journal of the Faculty of Medicine, University of Tabuk. All rights reserved with SMJS. © SMJS 2021

Variable	Moderate/high stressed mothers	P value	Moderate/high stressed fathers	P value
	(n=189)		(n=58)	
Living with their child /children				
Yes	186 (98.4%)	0.024*	56 (96.6%)	0.190
No	3 (1.6%)	0.024*	2 (3.4%)	0.180
Age				
≤29	35 (18.8%)		5 (8.6%)	
30-39	58 (30.7%)	0.016*	16 (27.6%)	0 111
40-49	81 (42.9%)	0.016*	25 (43.1%)	0.111
≥50	15 (7.9%)		12 (20.7%)	
Nationality				
Saudi	171 (90.5%)	0.020*	47 (81%)	0.042*
Non-Saudi	18 (9.5%)	0.039*	11 (19%)	0.045**
Educational level				
Less than high school	13 (6.9%)		9 (15.5%)	
High school/diploma	37 (19.6%)	0.270	12 (20.7%)	0.120
Collage or above	139 (73.5%)		37 (63.8%)	
Marital status				
Married	174 (92.1%)		56 (96.6%)	
		0.41		0.445
Divorced/ Widow	15 (7.9%)		2 (3.4%)	
Occupation				
Health care provider	14 (7.4%)		5 (8.6%)	
Retired	8 (4.2%)		6 (10.3%)	
Field work	43 (22.8%)	0.111	18 (31%)	0.586
Office work	39 (20.6%)		27 (46.6%)	
Not working/student	85 (45%)		2 (3.4%)	

Table	5	Association	between	socio-	demogra	ohics o	of the	parents	and	stress	levels	(n=247))
												`	/

* Statical significance p-value was detected at (p≤0.05) using Pearson chi-square test.

and psychological changes among their children. This finding is consistent with another study published in August 2020, which revealed no significant relationship when they used a linear regression model (R=.11, p=.31). [29] On the other hand, some studies conflict with our results. [30,31] Regarding demographical relation, our study revealed а statistically significant relationship between changes in the children's psychology and family history of COVID-19 infection (P=0.001). Additionally, the fathers' residence area was significantly associated with moderate to high stress levels (P=0.002). It could be understood as living in Makkah city was more challenging than living in Jeddah due to some specific precursory masseurs that have been done during the COVID-19 crisis. Also, parents' nationality showed a statical significance with having moderate/high levels of stress among parents (P=0.039 and 0.043, respectively, for mothers and fathers). Lastly, mothers who were 'living with their child' and also their 'age' had recorded a significant relation among mothers regarding having moderate to a high level of stress (P=0.024 and 0.016). Most of these results

SMJS is the official journal of the Faculty of Medicine, University of Tabuk. All rights reserved with SMJS. © SMJS 2021

are in agreement with other recorded studies in the literature. [29,30]

Limitations, Strengths, and Recommendations

The present study had some limitations that must be considered. First, most of the respondents in our study were married, Saudis, and highly educated. Therefore, the results may not apply to different family structures. Secondly, there is no validated tool (until the time of conducting the study) to assess changes in children's emotional state and behaviors during the COVID-19 pandemic. We adapted and revised a previously published tool from another study done in Italy and Spain. [7] Lastly, Sharedmethod bias is probable since parents are the only reporters for all study variables, and a self-report survey was the only tool used for data collection. Despite the limitations, our study's findings emphasize the importance of early detection of children's psychological changes as it may protect them from further complications. Our study results also reveal the large number of psychological stresses recorded in parents, which requires speedy finding effective solutions by individuals or institutions. Other researchers may find these results useful as base data for further investigations. However, we recommend more studies to be carried out regarding this issue in much better ways, avoiding this study's limitations.

CONCLUSION

The current study revealed no significant relationship between parental stress and psychological impact on their children. However, as an impact of COVID-19

quarantine, moderate to high-stress levels were seen in parents who live in Makkah. Additionally, significant changes in children's emotional states and behaviors were detected. We emphasize the importance of early detection of children's psychological changes to protect them from further complications. Ultimately, we recommend more studies to be carried out regarding this issue to clarify the finding in much better ways, avoiding the limitations of this study, as well as to investigate more the psychological effect of the crisis during this period with the different social and cultural changes on family's life, with the assessment of the level of needs for any phycological rehabilitation program for children and families.

ACKNOWLEDGMENTS Conflicts of Interest

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Authors' contribution

All authors have contributed equally in conceiving and designing the analysis, collecting the data, performing the analysis, and writing the manuscript of the paper.

Funding

The author received no financial support for the research, authorship, and/or publication of this article.

Availability of data and materials

The authors confirm that the data supporting the findings of this study are available and

SMJS is the official journal of the Faculty of Medicine, University of Tabuk. All rights reserved with SMJS. © SMJS 2021

will be provided by the corresponding author on a reasonable request.

Disclaimer

All opinions shared in this article are those of the authors and do not necessarily reflect the official policy or position of any affiliated agency of the authors.

Abbreviations

COVID-19: Coronavirus disease 2019 WHO: World Health Organization CI: confidence interval IRB: Internal Review Board UQU: Umm Al-Qura University SPSS: Statistical Package for the Social Sciences PCC: Pearson's Correlation Coefficient PSS: Perceived Stress Score

REFERENCES

1. Saudi Center for Disease Prevention and Control. Coronavirus Disease 19 (COVID-19) Guidelines. February 2020.

2. Ministry of Health. Novel Coronavirus (COVID-19) [Internet]. June 17 2020. [cited 2020 August 27]. Available from: https://www.moh.gov.sa/en/HealthAwareness/Educat ionalContent/Corona/Pages/corona.aspx

3. World Health Organization. WHO Coronavirus Disease (COVID-19) Dashboard. August 26 2020.

4. Ebrahim SH, Ahmed QA, Gozzer E, Schlagenhauf P, Memish ZA. Covid-19 and community mitigation strategies in a pandemic. Vol. 368, BMJ (Clinical research ed.). England; 2020. p. m1066.

5. Al-Tawfiq JA, Memish ZA. COVID-19 in the Eastern Mediterranean Region and Saudi Arabia: prevention and therapeutic strategies. Int J Antimicrob Agents [Internet]. 2020/04/04. 2020

May;55(5):105968. Available from: https://pubmed.ncbi.nlm.nih.gov/32259576

6. Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N, et al. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. Lancet. 2020;

7. Orgilés M, Morales A, Delvecchio E, Mazzeschi C, Espada JP. Immediate psychological effects of the COVID-19 quarantine in youth from Italy and Spain. 2020;

8. Centers for Disease Control and Prevention. Coronavirus Disease 2019 (COVID-19)- Helping Children Cope. July 1, 2020.

9. Dalton L, Rapa E, Stein A. Protecting the psychological health of children through effective communication about COVID-19. Lancet Child Adolesc Heal. 2020;4(5):346–7.

10. Jiao WY, Wang LN, Liu J, Fang SF, Jiao FY, Pettoello-Mantovani M, et al. Behavioral and emotional disorders in children during the COVID-19 epidemic. J Pediatr. 2020;221:264.

11. Goldschmidt K. The COVID-19 pandemic: Technology use to support the well-being of children. J Pediatr Nurs. 2020;

12. Davico C, Ghiggia A, Marcotulli D, Ricci F, Amianto F, Vitiello B. Psychological Impact of the COVID-19 Pandemic on Adults and Their Children in Italy. Available SSRN 3576933. 2020;

13. Sher L. COVID-19, anxiety, sleep disturbances and suicide. Sleep Med. 2020;

14. Chung SKG, Lanier P, Wong P. Mediating Effects of Parental Stress on Harsh Parenting And Parent-child Relationship During Coronavirus (COVID-19) Pandemic in Singapore. 2020;

15. Dean AG SK SM. Open Source Epidemiologic Statistics for Public Health [Internet]. April 6, 2013. [cited 2020 August 27]. Available from: https://www.openepi.com/SampleSize/SSPropor.htm

SMJS is the official journal of the Faculty of Medicine, University of Tabuk. All rights reserved with SMJS. © SMJS 2021

16. Cohen S, Kamarck T, Mermelstein R. A Global Measure of Perceived Stress. J Health Soc Behav [Internet]. 1983 August 17;24(4):385–96. Available from: http://www.jstor.org/stable/2136404

17. Ministry of Health. Covid19 Command and Control Center CCC. Official Updates Coronavirus -COVID-19 in Saudi Arabia [Internet]. 2020. [cited 2020 August 27]. Available from: https://covid19.moh.gov.sa

18. Liu JJ, Bao Y, Huang X, Shi J, Lu L. Mental health considerations for children quarantined because of COVID-19. Lancet Child Adolesc Heal [Internet]. 2020/03/27. 2020 May;4(5):347–9. Available from: https://pubmed.ncbi.nlm.nih.gov/32224303

19. Wang G, Zhang Y, Zhao J, Zhang J, Jiang F. Mitigate the effects of home confinement on children during the COVID-19 outbreak. Lancet (London, England) [Internet]. 2020/03/04. 2020 March 21;395(10228):945–7. Available from: https://pubmed.ncbi.nlm.nih.gov/32145186

20. Sprang G, Silman M. Posttraumatic stress disorder in parents and youth after health-related disasters. Disaster Med Public Health Prep. 2013 Feb;7(1):105–10.

21. World Health Organization. Coronavirus disease (COVID-19) advice for the public [Internet]. June 4 2020. [cited 2020 August 27]. Available from: https://www.who.int/emergencies/diseases/novel-coronavirus-2019/advice-for-public

22. Brown SM, Doom JR, Lechuga-Peña S, Watamura SE, Koppels T. Stress and parenting during the global COVID-19 pandemic. Child Abuse Negl [Internet]. 2020;104699. Available from: http://www.sciencedirect.com/science/article/pii/S01 45213420303549

23. Griffith AK. Parental Burnout and Child Maltreatment During the COVID-19 Pandemic. J Fam Violence [Internet]. 2020; Available from: https://doi.org/10.1007/s10896-020-00172-2

24. Yuan R, Xu Q, Xia C, Lou C, Xie Z, Ge Q, et al. Psychological status of parents of hospitalized children during the COVID-19 epidemic in China. Psychiatry Res [Internet]. 2020;288:112953. Available from: http://www.sciencedirect.com/science/article/pii/S01 65178120305539

25. Wu M, Xu W, Yao Y, Zhang L, Guo L, Fan J, et al. Mental health status of students' parents during COVID-19 pandemic and its influence factors. Gen Psychiatry [Internet]. 2020 Jul 21;33(4):e100250. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7387 315/

26. Spinelli M, Lionetti F, Pastore M, Fasolo M. Parents' Stress and Children's Psychological Problems in Families Facing the COVID-19 Outbreak in Italy. Front Psychol [Internet]. 2020;11:1713. Available from:

https://www.frontiersin.org/article/10.3389/fpsyg.202 0.01713

27. Jeong H, Yim HW, Song Y-J, Ki M, Min J-A, Cho J, et al. Mental health status of people isolated due to Middle East Respiratory Syndrome. Epidemiol Health. 2016;38:e2016048.

28. Dubey S, Biswas P, Ghosh R, Chatterjee S, Dubey MJ, Chatterjee S, et al. Psychosocial impact of COVID-19. Diabetes Metab Syndr [Internet]. 2020 May 27;14(5):779–88. Available from: https://pubmed.ncbi.nlm.nih.gov/32526627

29. Shorer M, Leibovich L. Young children's emotional stress reactions during the COVID-19 outbreak and their associations with parental emotion regulation and parental playfulness. Early Child Dev Care [Internet]. 2020 August 20;1–11. Available from: https://doi.org/10.1080/03004430.2020.1806830

30. Romero-González M, Marín E, Guzmán-Parra J, Navas P, Aguilar JM, Lara JP, et al. [Relationship between parental stress and psychological distress and emotional and behavioural problems in pre-school children with autistic spectrum disorder]. An Pediatr

SMJS is the official journal of the Faculty of Medicine, University of Tabuk. All rights reserved with SMJS. © SMJS 2021

(Barc) [Internet]. 2020; Available from: https://doi.org/10.1016/j.anpedi.2020.03.012

31. Goodman R. The Strengths and Difficulties Questionnaire: a research note. J Child Psychol Psychiatry. 1997 Jul;38(5):581–6.



To receive the weekly newsletter of the Faculty of Medicine, University of Tabuk, KSA, please send your email to mededutabuk@ut.edu.sa

SMJS is the official journal of the Faculty of Medicine, University of Tabuk. All rights reserved with SMJS. © SMJS 2021